arXiv:1201.3818v1 [math.CO] 18 Jan 2012

# A note on heterochromatic cycles of length 4 in edge-colored graphs[*]

Bo Ning and Shenggui Zhang[†]

Department of Applied Mathematics, Northwestern Polytechnical University,
Xi'an, Shaanxi 710072, P.R. China

## Abstract

Let $G$ be an edge-colored graph. A heterochromatic cycle of $G$ is one in which every two edges have different colors. For a vertex $v \in V(G)$, let $CN(v)$ denote the set of colors which are assigned to the edges incident to $v$. In this note we prove that $G$ contains a heterochromatic cycle of length 4 if $G$ has $n \geq 60$ vertices and $|CN(u) \cup CN(v)| \geq n-1$ for every pair of vertices $u$ and $v$ of $G$. This extends a result of Broersma et al. on the existence of heterochromatic cycles of length 3 or 4.

**Keywords:** edge-colored graphs; heterochromatic cycles

## 1 Introduction

We use Bondy and Murty [2] for terminology and notation not defined here and consider finite simple graphs only.

Let $G = (V, E)$ be a graph, $H$ a subgraph, and $v$ a vertex of $G$. We use $N_H(v)$ to denote the set, and $d_H(v)$ the number, of neighbors of $v$ in $H$, and call $d_H(v)$ the *degree* of $v$ in $H$. An *edge-coloring* of $G$ is a function $C : E \longrightarrow \mathbb{N}^+$, where $\mathbb{N}^+$ is the set of positive integers. We call $G$ an *edge-colored graph* if it is assigned such a coloring $C$, and denote this edge-colored graph by $(G, C)$. We use $CN_H(v)$ to denote the set of colors of the edges joining $v$ and its neighbors in $H$ when $v \notin V(H)$ and those incident to $v$ in $H$ when $v \in V(H)$, and call $d^c_H(v) = |CN_H(v)|$ the *color degree* of $v$ in $H$. When no confusion occurs, we use $G$ instead of $(G, C)$, and $N(v)$, $d(v)$, $CN(v)$ and $d^c(v)$ instead of $N_G(v)$, $d_G(v)$, $CN_G(v)$ and $d^c_G(v)$, respectively.

---

[*] Supported by NSFC (No. 10871158).

[†] Corresponding author. E-mail address: sgzhang@nwpu.edu.cn (S. Zhang).

A subgraph $H$ of an edge-colored graph $G$ is called *heterochromatic* if every two of its edges have different colors. Recently, heterochromatic subgraphs in edge-colored graphs have received much attentions. Heterochromatic matchings were studied in [9, 11, 12]. Chen and Li [5, 6] studied long heterochromatic paths in edge-colored graphs. Albert et al. [1] studied heterochromatic Hamilton cycles in edge-colored complete graphs. Heterochromatic cycles of small lengths were studied in [4, 10, 13, 14]. For a survey on the study of heterochromatic subgraphs in edge-colored graphs, we refer to [8].

In the following we use $C_k$ to denote a cycle of length $k$. Broersma et al. [4] considered the existence of short heterochromatic cycles in edge-colored graphs and got the following result.

**Theorem 1** (Broersma, Li, Woeginger and Zhang [4])**.** *Let $G$ be an edge-colored graph of order $n \geq 4$ such that $|CN(u) \cup CN(v)| \geq n-1$ for every pair of vertices $u$ and $v$ of $G$. Then $G$ contains a heterochromatic $C_3$ or a heterochromatic $C_4$ .*

Li and Wang [10] gave a result on the existence of a heterochromatic $C_3$ or a heterochromatic $C_4$ under the color degree condition. At the same time, they obtained a result on the existence of a heterochromatic $C_3$ in edge-colored graphs.

**Theorem 2** (Li and Wang [10])**.** *Let $G$ be an edge-colored graph of order $n \geq 3$. If $d^c(v) \geq (\frac{4\sqrt{7}}{7} - 1)n + 3 - \frac{4\sqrt{7}}{7}$ for every vertex $v \in V(G)$, then $G$ contains either a heterochromatic $C_3$ or a heterochromatic $C_4$.*

**Theorem 3** (Li and Wang [10])**.** *Let $G$ be an edge-colored graph of order $n \geq 3$. If $d^c(v) \geq \frac{\sqrt{7}+1}{6}n$ for every vertex $v \in V(G)$, then $G$ contains a heterochromatic $C_3$.*

Later, Wang et al. [13] got a result on the existence of a heterochromatic $C_4$ in edge-colored triangle-free graphs. Zhu [14] further extended this result to edge-colored bipartite graphs.

**Theorem 4** (Wang, Li, Zhu and Liu [13])**.** *Let $G$ be an edge-colored triangle-free graph of order $n \geq 9$. If $d^c(v) \geq \frac{3-\sqrt{5}}{2}n + 1$ for every vertex $v \in V(G)$, then $G$ contains a heterochromatic $C_4$.*

**Theorem 5** (Zhu [14])**.** *Let $G$ be an edge-colored bipartite graph of order $n \geq 6$. If $d^c(v) \geq \frac{(\sqrt{5}-1)n}{4} + 1$ for every vertex $v \in V(G)$, then $G$ contains a heterochromatic $C_4$.*

In this note we prove the following result which is an extension of Theorem 1.

**Theorem 6.** *Let $G$ be an edge-colored graph of order $n \geq 60$ such that $|CN(u) \cup CN(v)| \geq n-1$ for every pair of vertices $u$ and $v$ of $G$. Then $G$ contains a heterochromatic $C_4$.*

We can not provide examples to show that the restriction $n \geq 60$ can be reduced and the condition $|CN(u) \cup CN(v)| \geq n-1$ can be weakened in Theorem 6. However, we can show that the lower bound of $|CN(u) \cup CN(v)|$ must be more than $\sqrt{2n-3}$ in order to guarantee the existence of a heterochromatic $C_4$.

A finite projective plane $\mathcal{P}$ is a pair of sets $(P, L)$ where $P$ is a set of points and $L$ is a set of lines such that

$(i)$ any two points in $P$ lie on only one line in $L$;
$(ii)$ any two lines in $L$ meet in only one point in $P$; and
$(iii)$ there are four points in $P$ no three of which lie on a line in $L$.

It is proved that (see [3], Exercise 1.3.13) for any integer $t$, there exists a finite projective plane $(P, L)$ such that each point in $P$ lies on $t+1$ lines in $L$, each line in $L$ contains $t+1$ points in $P$, and $|P| = |L| = t^2 + t + 1$.

Let $\mathcal{P}$ be a finite projective plane with $|P| = |L| = t$. We define the *incidence graph* $I(\mathcal{P})$ of $\mathcal{P}$ as the bipartite graph with bipartition $(P, L)$ and edges $pl$ when $p \in P$ lies on $l \in L$ in $\mathcal{P}$. Let $u, v$ be two vertices of $I(\mathcal{P})$. If $u, v \in P$ or $u, v \in L$, then $|N(u) \cup N(v)| = 2t+1$. If one of them (say $u$) is in $P$ and the other ($v$) is in $L$, then $|N(u) \cup N(v)| = 2t+1$ when $u$ lies on $v$ and $|N(u) \cup N(v)| = 2t+2$ otherwise. Therefore, we have $|N(u) \cup N(v)| \geq 2t+1 = \sqrt{2n-3}$. On the other hand, it is easy to see that $I(\mathcal{P})$ contains no cycles of length 4. Otherwise, there exist two points in $P$ which lie on two lines, contradicting to the definition of finite projective planes.

If we assign an edge coloring to $I(\mathcal{P})$ such that every two of its edges have different colors, then we can get the following proposition.

**Proposition 1.** *Let $t$ be an integer. Then there exists an edge-colored graph $G$ of order $n = 2(t^2 + t + 1)$ such that $|CN(u) \cup CN(v)| \geq \sqrt{2n-3}$ for every pair of vertices $u$ and $v$ of $G$ and $G$ contains no heterochromatic cycles of length 4.*

## 2 Proof of Theorem 6

We first give a lemma, which will be used in the proof of Theorem 6.

**Lemma 1.** *Let $G$ be an edge-colored graph. Then $G$ contains a spanning bipartite subgraph $H$ such that $2d^c_H(v) + 3d_H(v) \geq d^c_G(v) + d_G(v)$ for every vertex $v \in V(H)$.*

*Proof.* We choose a spanning bipartite subgraph $H$ of $G$ such that $f(H) = |E(H)| + \sum_{v \in V(H)} d^c_H(v)$ is as large as possible and show that $2d^c_H(v) + 3d_H(v) \geq d^c_G(v) + d_G(v)$ for every vertex $v \in V(H)$.

Assume that the bipartition of $H$ is $(X, Y)$. Then any edge $xy$ of $G$ with $x \in X$ and $y \in Y$ is also an edge of $H$. Otherwise, we have $f(H + xy) > f(H)$, contradicting to the choice of $H$. It can be seen that $d^c_H(x) = |CN_{G[Y]}(x)|$ for $x \in X$ and $d^c_H(y) = |CN_{G[X]}(y)|$ for $y \in Y$.

Suppose that there exists a vertex $w \in V(H)$ such that

$$2d^c_H(w) + 3d_H(w) < d^c_G(w) + d_G(w). \tag{1}$$

Without loss of generality, we assume $w \in X$. Let $H'$ be the spanning bipartite subgraph of $G$ with bipartition $(X\backslash\{w\}, Y \cup \{w\})$ and edge set $E(H) \cup \{wx : x \in X \setminus \{w\}\} \setminus \{wy : y \in Y\}$. Then, we have

$$|E(H')| - |E(H)| = (d_G(w) - d_H(w)) - d_H(w) = d_G(w) - 2d_H(w). \tag{2}$$

On the other hand, noting that

$$\begin{aligned}
d^c_{H'}(w) - d^c_H(w) &= |CN_{G[X]}(w)| - |CN_{G[Y]}(w)| \\
&= |CN_{G\backslash G[Y]}(w)| - |CN_{G[Y]}(w)| \\
&\geq |CN_G(w)| - 2|CN_{G[Y]}(w)| \\
&= d^c_G(w) - 2d^c_H(w)
\end{aligned}$$

and

$$\begin{aligned}
\sum_{v \in V\backslash\{w\}} (d^c_{H'}(v) - d^c_H(v)) &= \sum_{v \in X\backslash\{w\}} (d^c_{H'}(v) - d^c_H(v)) + \sum_{v \in Y} (d^c_{H'}(v) - d^c_H(v)) \\
&\geq \sum_{v \in Y} (d^c_{H'}(v) - d^c_H(v)) \\
&= \sum_{v \in Y} (|CN_{G[X\backslash\{w\}]}(v)| - |CN_{G[X]}(v)|) \\
&\geq -\sum_{v \in Y} |CN_{G[\{w\}]}(v)| \\
&= -d_H(w),
\end{aligned}$$

we have

$$\begin{aligned}
\sum_{v \in V} d^c_{H'}(v) - \sum_{v \in V} d^c_H(v) &= \sum_{v \in V\backslash\{w\}} (d^c_{H'}(v) - d^c_H(v)) + (d^c_{H'}(w) - d^c_H(w)) \\
&\geq (d^c_G(w) - 2d^c_H(w)) - d_H(w).
\end{aligned}$$

That is,

$$\sum_{v \in V} d^c_{H'}(v) - \sum_{v \in V} d^c_H(v) \geq d^c_G(w) - 2d^c_H(w) - d_H(w). \tag{3}$$

By (1), (2) and (3), we get

$$f(H') - f(H) \geq d_G(w) + d^c_G(w) - 2d^c_H(w) - 3d_H(w) > 0,$$

which contradicts to the choice of $H$.

The proof is complete. $\square$

**Proof of Theorem 6**

We denote $\delta^c(G) = \min\{d^c(v) : v \in V(G)\}$ and distinguish two cases.

**Case 1.** $\delta^c(G) = n - 1$.

Assume that $G$ contains no heterochromatic cycles of length 4. Then we have $d^c(v) = d(v) = n - 1$ for every vertex $v \in V(G)$. Hence $G$ is complete and $C(vx) \neq C(vy)$ for every vertex $v \in V(G)$, where $x, y \in N(v)$.

**Claim 1.** Let $x_1, x_2, x_3, x_4$ be four vertices of $V(G)$. If $C(x_1x_2) \neq C(x_3x_4)$, then $C(x_2x_3) = C(x_1x_4)$.

*Proof.* Suppose that $C(x_2x_3) \neq C(x_1x_4)$. From the above discussion, we know that $C(x_1x_2) \neq C(x_3x_4)$, $C(x_1x_2) \neq C(x_2x_3)$, $C(x_2x_3) \neq C(x_3x_4)$, $C(x_3x_4) \neq C(x_4x_1)$ and $C(x_4x_1) \neq C(x_1x_2)$. With $C(x_2x_3) \neq C(x_1x_4)$, we see that $x_1x_2x_3x_4x_1$ is a heterochromatic $C_4$, a contradiction. $\square$

Let $y_1, y_2, y_3, y_4, y_5$ be five vertices of $V(G)$. Without loss of generality, we assume $C(y_5y_k) = k$ for $k \in \{1, 2, 3, 4\}$. Since $C(y_1y_2) \neq C(y_1y_5)$ and $C(y_1y_2) \neq C(y_2y_5)$, we have $C(y_1y_2) \neq 1$ and $C(y_1y_2) \neq 2$. Hence $C(y_1y_2) \geq 3$.

Suppose that $C(y_1y_2) = 3$. Since $C(y_1y_2) \neq C(y_4y_5)$, it follows from Claim 1 that $C(y_2y_4) = C(y_5y_1)$. With $C(y_3y_5) \neq C(y_1y_5)$, we have $C(y_2y_4) \neq C(y_3y_5)$. Then by Claim 1, we obtain that $C(y_3y_4) = C(y_2y_5)$. At the same time, by $C(y_1y_2) \neq C(y_4y_5)$ and Claim 1, we have $C(y_1y_4) = C(y_2y_5)$. Hence $C(y_1y_4) = C(y_3y_4)$, a contradiction.

Suppose that $C(y_1y_2) = 4$. Since $C(y_1y_2) \neq C(y_3y_5)$, it follows from Claim 1 that $C(y_2y_3) = C(y_1y_5)$. With $C(y_4y_5) \neq C(y_1y_5)$, we get $C(y_2y_3) \neq C(y_4y_5)$. Then by Claim 1, we obtain that $C(y_3y_4) = C(y_2y_5)$. Similarly, with $C(y_1y_2) \neq C(y_2y_5)$, we have $C(y_1y_2) \neq C(y_3y_4)$. Then by Claim 1, we have $C(y_1y_4) = C(y_2y_3)$. Hence, $C(y_1y_4) = C(y_1y_5)$, a contradiction.

Suppose that $C(y_1y_2) \geq 5$. Since $C(y_1y_2) \neq C(y_4y_5)$, it follows from Claim 1 that $C(y_2y_4) = C(y_1y_5)$. At the same time, by $C(y_1y_2) \neq C(y_3y_5)$ and Claim 1, we get $C(y_2y_3) = C(y_1y_5)$. Hence $C(y_2y_4) = C(y_2y_3)$, a contradiction.

The proof of Case 1 is complete.

**Case 2.** $\delta^c(G) \le n-2$.

Let $w$ be a vertex with $d^c(w) = \delta^c(G)$ and denote $\delta^c(G) = k$. Let $T$ be a subset of $N(w)$ such that $|T| = k$ and $C(wx) \neq C(wy)$ for every two vertices $x, y \in T$. Without loss of generality, set $T(w) = \{x_1, x_2, \ldots, x_k\}$ and assume that $C(wx_i) = i$ for $i \in \{1, 2, \ldots, k\}$. Set $G_1 = G[T \cup \{w\}]$ and $G_2 = G[V(G) \setminus V(G_1)]$. Since $|V(G_1)| = k+1 \le n-1$, we have $V(G_2) \neq \emptyset$.

**Case 2.1.** There exits a vertex $z \in V(G_2)$ such that $|CN_{G_1}(z) \setminus CN(w)| \ge 2$.

By the choice of $T$, if $v$ is a neighbor of $z$ such that $C(vz) \in CN_{G_1}(z) \setminus CN(w)$, then $v \neq w$. Since $|CN_{G_1}(z) \setminus CN(w)| \ge 2$, we can choose $x_s, x_t \in T$ with $\{C(x_s z), C(x_t z)\} \subseteq CN_{G_1}(z) \setminus CN(w)$. Obviously, $wx_szx_tw$ is a heterochromatic $C_4$.

**Case 2.2.** $|CN_{G_1}(v) \setminus CN(w)| \le 1$ for every vertex $v \in V(G_2)$.

**Claim 2.** $|CN_{G_2}(v)| = |V(G_2)| - 1$ for every vertex $v \in V(G_2)$.

*Proof.* First, we have $|CN_{G_1}(v) \backslash CN(w)| \le 1$. If follows from $|CN(w)| = k$ that $|CN(w) \cup CN_{G_1}(v)| \le k+1$. Note that $|CN(w) \cup CN(v)| \ge n-1$, we have $|CN(v) \backslash CN_{G_1}(v)| \ge n-k-2$. On the other hand, we have $|CN(v) \backslash CN_{G_1}(v)| \le |CN_{G_2}(v)| \le d_{G_2}(v) \le |V(G_2)| - 1 = n-k-2$. Thus, $|CN_{G_2}(v)| = |V(G_2)| - 1$, where $|V(G_2)| = n-k-1$. □

**Case 2.2.1.** $k \le n-6$.

Note that $|V(G_2)| = n-k-1 \ge 5$. Then as in Case 1, we can prove that there exists a heterochromatic $C_4$ in $G_2$, which is also a heterochromatic $C_4$ in $G$.

**Case 2.2.2.** $k \ge n-5$.

By Lemma 1, there is a spanning bipartite subgraph $H$ such that

$$2d_H^c(v) + 3d_H(v) \ge d_G^c(v) + d_G(v) \tag{4}$$

for every vertex $v \in V(H)$. It is not difficult to see that

$$d_H(v) - d_H^c(v) \le d_G(v) - d_G^c(v) \tag{5}$$

and

$$d_G(v) - d_G^c(v) \le d_G(v) - \delta^c(G) \le (n-1) - (n-5) = 4. \tag{6}$$

Together with (5) and (6), we have

$$d^c_H(v) - d_H(v) \geq -4. \tag{7}$$

Then, combining (4) with (7), we obtain

$$d^c_H(v) \geq \frac{1}{5}(d^c_G(v) + d_G(v) - 12) \geq \frac{2n-22}{5} > \frac{(\sqrt{5}-1)n}{4} + 1$$

when $n \geq 60$. By Theorem 5, there is a heterochromatic $C_4$ in $H$, which is also a heterochromatic $C_4$ in $G$.

The proof is complete. $\square$

## 3 Remarks

Our proof of Theorem 6 mainly relies on Lemma 1 and Theorem 5.

Lemma 1 was motivated by the following result due to Erdös [7].

**Theorem 7** (Erdös [7])**.** *Let $G$ be a graph. Then $G$ contains a spanning bipartite subgraph $H$ such that $d_H(v) \geq \frac{1}{2}d_G(v)$ for all $v \in V(H)$.*

We have the following problem.

**Problem 1.** Let $G$ be an edge-colored graph. Does $G$ contain a spanning bipartite subgraph $H$ such that $d^c_H(v) \geq \frac{1}{2}d^c_G(v)$?

Zhu [13] pointed out that if the answer to the following conjecture is positive, then it would result in an improvement to Theorem 5.

**Conjecture 1** ([11])**.** Let $D$ be a directed bipartite graph with bipartition $(A, B)$. If $d^+(u) > \frac{|B|}{3}$ for $u \in A$ and $d^+(v) \geq \frac{|A|}{3}$ for $v \in B$, or $d^+(u) \geq \frac{|B|}{3}$ for $u \in A$ and $d^+(v) > \frac{|A|}{3}$ for $v \in B$, then there exists a directed $C_4$ in $D$.

From our proof of Theorem 6, we know that if the answer to Problem 1 or Conjecture 1 is positive, then the restriction $n \geq 60$ in Theorem 6 can be reduced.

## References

[1] M. Albert, A. Frieze and B. Reed, Multicolored Hamilton cycles, *Electron. J. Comb.* **2** (1995), #R10.

[2] J.A. Bondy and U.S.R. Murty, Graph Theory with Applications, Macmillan London and Elsevier, New York (1976).

[3] J.A. Bondy and U.S.R. Murty, Graph Theory, GTM-244, Springer, Berlin (2008).

[4] H.J. Broersma, X. Li, G. Woegingerr and S. Zhang, Paths and cycles in colored graphs, *Australas. J. Comb.* **31** (2005), 297-305.

[5] L. Chen and X. Li, long heterochromatic paths in edge-colored graphs, *Electron. J. Comb.* **12** (2005), #R33.

[6] L. Chen and X. Li, Color neighborhood union conditions for long heterochromatic paths in edge-colored graphs, *Electron. J. Comb.* **14** (2007), #R77.

[7] P. Erdös, On some extremal problem in graph theory, *Isral. J. Math.* **3** (1965), 113-116.

[8] M. Kano and X. Li, Monochromatic and heterochromatic subgraphs in edge-colored graphs-a survey, *Graphs Combin.* **93** (2009), 129-139.

[9] T.D. Lesaulnier, C. Stocker, P.S. Wenger and D.B. West, Rainbow matching in edge-colored graphs, *Electron. J. Comb.* **17** (2010), #N26.

[10] H. Li and G. Wang, Color degree and heterochromatic cycles in edge-colored graphs, *European J. Combin.*, to appear.

[11] G. Wang, Rainbow matchings in properly edge colored graphs, *Electron. J. Comb.* **18** (2011), #P162.

[12] G. Wang and H. Li, Heterochromatic matchings in edge-colored graphs, *Electron. J. Comb.* **15** (2008), #R138.

[13] G. Wang, H. Li, Y. Zhu and G. Liu, A note on heterochromatic $C_4$ in edge-colored triangle-free graphs, *Graphs Combin.*, to appear.

[14] Yan Zhu, Randić Index and Cycles in Graphs and Applications, PhD thesis, Paris-Sud 11 University (France), Shandong University (China), (2010).